

- a new cause of disease? *Am J Med Genet* 91: 22–8
- Bianchi DW, Zickwolf GK, Weil GJ, Sylvester S, DeMaria MA (1996) Male fetal progenitor cells persist in maternal blood for as long as 27 years postpartum. *Proc Natl Acad Sci USA* 93:705–8
- Gannage M, Amoura Z, Lantz O, Piette JC, Caillat-Zucman S (2002) Feto-maternal microchimerism in connective tissue diseases. *Eur J Immunol* 32:3405–13
- Grove JE, Bruscia E, Krause DS (2004) Plasticity of bone marrow-derived stem cells. *Stem Cells* 22:487–500
- Johnson KL, Bianchi DW (2004) Fetal cells in maternal tissue following pregnancy: what are the consequences? *Hum Reprod Update* 10:497–502
- Khosrotehrani K *et al.* (2006) Presence of chimeric maternally-derived keratinocytes in cutaneous inflammatory diseases of children: the example of pityriasis lichenoides. *J Invest Dermatol* 126:345–348
- Lo YM *et al.* (1996) Two-way cell traffic between mother and fetus: biologic and clinical implications. *Blood* 88:4390–5
- Regauer S, Schwaiger P, Liegl B, Klitschar M (2004) Fetal microchimerism is common in normal and diseased vulvar skin. *J Invest Dermatol* 122:1059–60
- Starzl TE (2004) Chimerism and tolerance in transplantation. *Proc Natl Acad Sci USA* 101(Suppl 2):14607–14

See related article on page 374

## (P)PARsing Epidermal Development

Sandrine Dubrac<sup>1</sup> and Matthias Schmuth<sup>1</sup>

Overexpression of PPAR- $\alpha$ , a developmental transcription factor important in epidermal embryogenesis, in basal keratinocytes causes epidermal thinning when activated constitutively during development, but not if activated in adults; and lack of PPAR- $\alpha$  transiently delays stratum corneum formation within a window late in epidermal development (day 18.5 to birth). In contrast, pharmacologic activation of PPAR- $\alpha$  inhibits proliferation and induces differentiation in mouse epidermis regardless of developmental stage. Thus, PPAR- $\alpha$  is an important regulator of epidermal homeostasis.

*Journal of Investigative Dermatology* (2006) 126, 241–242. doi:10.1038/sj.jid.5700137

Nuclear hormone receptors are transcription factors that regulate the expression of target genes by binding to regulatory DNA sequences and interacting with co-regulatory protein complexes. A large molecular family of these receptors has been identified by homology searches. Soon after the sequences became available, individual receptors were characterized for their tissue distribution, ligand identity, patterns of target gene activation, and interaction with co-regulatory proteins. To better understand their physiological roles, individual receptors were tested in genetic animal models of loss and gain of function.

In this fashion, peroxisome proliferator-activated receptor- $\alpha$  (PPAR $\alpha$ ) was found to be a nuclear hormone receptor that is activated by fatty acid-derived

ligands. Since Issemann and Green cloned mouse PPAR $\alpha$  in 1990 and Sher *et al.* cloned its human homologue in 1993 (Issemann and Green, 1990; Sher *et al.*, 1993), several groups have generated mouse models of PPAR $\alpha$  deficiency. Such animals displayed abnormal lipid and xenobiotic metabolism in the liver, heart, muscle, and kidney, indicating a role of PPAR $\alpha$  in fatty acid oxidation and detoxification of xenobiotic compounds. Although PPAR $\alpha$  was originally evaluated for its systemic activities, its expression was soon also noted in skin.

PPAR $\alpha$  is present in both epidermis and dermis beginning at day 13.5 of development. Yet shortly after birth it becomes undetectable in the interfollicular epidermis, although expression persists in the hair follicles (Michalik

*et al.*, 2001). Injury to adult murine skin, such as hair plucking, induces re-expression of PPAR $\alpha$  in the adult interfollicular epidermis, and re-expression can also be observed in the edges of full-thickness wounds. Conversely, in PPAR $\alpha$ -deficient mice, the early phase of wound healing is delayed, and this delay is retained when the deficiency is targeted to the epidermis only and not to the dermis (Michalik *et al.*, 2005). In pups lacking PPAR $\alpha$ , a delay in stratum corneum formation is observed between day 18.5 of epidermal development and birth (Schmuth *et al.*, 2002), whereas in PPAR $\alpha$ -deficient adults, only a modest decrease in the expression of involucrin, loricrin, and filaggrin persists. This indicates that other mediators can compensate for the absence of PPAR $\alpha$ ; that is, there is redundancy (Komuves *et al.*, 2000).

In this issue, Gonzalez *et al.* (2006) report on the skin phenotype of transgenic mice constitutively overexpressing PPAR $\alpha$  in the epidermis. These mice die within 2 days after birth, presumably because of abnormal development of the tongue and mammary gland epithelia; overexpression of PPAR $\alpha$  also results in epidermal thinning and sparse fur in these animals, which could contribute to the lethality. Importantly, corresponding to the transient effects of PPAR $\alpha$  deficiency on developing epidermis, PPAR $\alpha$  overexpression exerts its effects only during a developmental window; that is, after birth it does not cause the abnormalities.

Consequences of a gain of PPAR $\alpha$  function have previously been studied using pharmacologic activators. In explants of developing rat epidermis, the expression of proteins required for epidermal differentiation (filaggrin, loricrin, involucrin) was stimulated by the PPAR $\alpha$  agonist farnesol (Hanley *et al.*, 1997). In contrast to the VP16PPAR- $\alpha$  bitransgenic mice reported here (Gonzales *et al.*, 2006), there was a concomitant induction of the granular layer. These differences could be explained by differences between *in vitro* and *in vivo* experimental systems and by differences in the timing of the PPAR $\alpha$  signal. Nevertheless, in adult mice, pharmacologic activation of PPAR $\alpha$  induces epidermal differentiation, inhibits proliferation, and increases

<sup>1</sup>Department of Dermatology, Innsbruck Medical University, Innsbruck, Austria

Correspondence: Dr Sandrine Dubrac, Department of Dermatology, Innsbruck Medical University, Anichstraße 35, A6020 Innsbruck, Austria. Email: sandrine.dubrac@uibk.ac.at

keratinocyte death. These effects are absent in PPAR $\alpha$  knockout mice, which indicates that they are specifically mediated via PPAR $\alpha$  (Komuves *et al.*, 2000). Consistent with these findings, Gonzales *et al.* (2006) report here an increase in p21 (Waf1/Cip1) in cultured keratinocytes from 1-day-old neonate mice overexpressing PPAR $\alpha$ , which, by causing an arrest in the G1 phase of the cell cycle, may account for the coordinate regulation of proliferation, differentiation, and cell death.

How do these results complement our current knowledge on the transcriptional regulation of epidermal development? Aside from p21, which has been shown to act as a negative regulator of transcription, linking the Notch and Wnt signaling pathways in the control of keratinocyte growth (Devgan *et al.*, 2005), additional transcription factors have been implicated in skin morphogenesis. Whereas LEF-1 and TCF-3 have primarily been investigated for their role in the epidermal stem cell niche (Merrill *et al.*, 2001), the work of Gonzales *et al.* (2006) implies that PPAR $\alpha$  is involved in controlling the switch from proliferating basal to differentiating suprabasal keratinocytes. The precise compartmental roles of PPAR $\alpha$  remain to be determined, and it is certainly only one of multiple transcription factors involved in controlling epidermal transcription and is likely to interact with additional factors — p63, SP-1, the activator protein-1 transcription factor gene family, c-Myc, RelA, pRb, Klf4, and others (Dai and Segre, 2004). The challenge of future investigations in this field will be to delineate a more complete and possibly quantitative map of interactions among these factors to obtain a better understanding of how epidermal development is orchestrated.

Although it is clear that ligand binding controls the biological activity of PPAR $\alpha$ , the specific lipid ligands that PPAR $\alpha$

senses physiologically in keratinocytes remain unknown. There is a large number of potential long-chain unsaturated fatty acid ligands, including eicosanoids and leukotrienes, and it is likely that, physiologically, more than one lipid compound is involved. The levels of a variety of ligands, rather than the level of one specific lipid, may be important in determining the activity of the receptor during epidermal development. Because PPAR $\alpha$  regulates genes of lipid metabolism, a feedback loop is likely to exist. Additionally, some lipids may be potent activators whereas other lipids may actually behave as antagonists. The phenotype of the PPAR $\alpha$ -overexpressing mice generated by the Gonzalez group supports the notion of cross-talk between epidermal lipids and the proteins involved in epidermal development.

Because PPAR $\alpha$  can be activated by ligands, the results from the current report provide the rationale for the development of tailored drugs targeting this receptor. In addition to a role in epidermal proliferation and differentiation, PPAR $\alpha$  activation has also been shown to be anti-inflammatory, and the delay in wound healing in PPAR $\alpha$ -deficient mice has been ascribed to impaired neutrophil and monocyte recruitment during the initial inflammatory phase of wound healing (Michalik *et al.*, 2001). The combination of these effects could be beneficial in the treatment of inflammatory skin disease associated with disturbed epidermal homeostasis. Even though extrapolation from mouse to human is perilous, this research should stimulate translational studies in humans that may lead to new avenues of therapeutic intervention for common dermatologic skin disease.

#### ACKNOWLEDGMENT

We are grateful to Robert Kimmel (Stanford University) for critically reviewing the manuscript.

#### CONFLICT OF INTEREST

The authors state no conflict of interest.

#### REFERENCES

- Dai X, Segre JA (2004) Transcriptional control of epidermal specification and differentiation. *Curr Opin Genet Dev* 14:485–91
- Devgan V, Mammucari C, Millar SE, Briskin C, Dotto GP (2005) p21WAF1/Cip1 is a negative transcriptional regulator of Wnt4 expression downstream of Notch1 activation. *Genes Dev* 19:1485–95
- Gonzalez FJ, Yang Q, Yamada A, Kimura S, Peters JM (2006) Alterations in skin and stratified epithelia by constitutively activated PPAR $\alpha$ . *J Invest Dermatol* 126:374–385 ]
- Hanley K, Jiang Y, Crumrine D, Bass NM, Appel R, Elias PM *et al.* (1997) Activators of the nuclear hormone receptors PPARalpha and FXR accelerate the development of the fetal epidermal permeability barrier. *J Clin Invest* 100:705–12
- Issemann I, Green S (1990) Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347:645–50
- Komuves LG, Hanley K, Lefebvre AM, Man MQ, Ng DC, Bikle DD *et al.* (2000) Stimulation of PPARalpha promotes epidermal keratinocyte differentiation in vivo. *J Invest Dermatol* 115:353–60
- Merrill BJ, Gat U, DasGupta R, Fuchs E (2001) Tcf3 and Lef1 regulate lineage differentiation of multipotent stem cells in skin. *Genes Dev* 15:1688–705
- Michalik L, Desvergne B, Tan NS, Basu-Modak S, Escher P, Rieusset J *et al.* (2001) Impaired skin wound healing in peroxisome proliferator-activated receptor (PPAR)alpha and PPARbeta mutant mice. *J Cell Biol* 154:799–814
- Michalik L, Feige JN, Gelman L, Pedrazzini T, Keller H, Desvergne B *et al.* (2005) Selective expression of a dominant-negative form of peroxisome proliferator-activated receptor in keratinocytes leads to impaired epidermal healing. *Mol Endocrinol* 19:2335–48
- Schmuth M, Schoonjans K, Yu QC, Fluhr JW, Crumrine D, Hachem JP *et al.* (2002) Role of peroxisome proliferator-activated receptor alpha in epidermal development in utero. *J Invest Dermatol* 119:1298–303
- Sher T, Yi HF, McBride OW, Gonzalez FJ (1993) cDNA cloning, chromosomal mapping, and functional characterization of the human peroxisome proliferator activated receptor. *Biochemistry* 32:5598–604